Working Group 2, Task 2: Survey on Quality Assurance Procedures, 2012

Dr Phil Gilvin Public Health England, UK




Questions on QA Schemes

Box F: What quality system do you use? Multiple answers are possible Certified by third party ISO 9000 series Compliant Certified ISO 17025 Other Compliant Certified Internal procedure based on: RP 73 (EUR 14852) RP 160 ☑ IAEA Safety Standard Series RS-G-1.3 Other




Quality Systems to which IMSs are Certified





QA Systems with which IMSs claim Compliance





"Other" Answers

- 5 x National systems
- 3 x In-house systems
- 5 misunderstood the question:
 - ISO 17020
 - ISO 1757
 - IEC 61066



QA Systems Implemented

	2003	2012
ISO 9000 Series	18%	41%
ISO 17025	24%	68%
Any formal QA	Up to 83%	91%

Note: multiple answers possible. Includes national QA schemes. 2012 figures include both those IMSs with formal certification and those who claim their system is compliant.



Questions on QA Schemes

Box G: What quality audits you undergo? Multiple answers are possible				
	Please give frequency of audits	Interval		
External - form	nal quality standard e.g. ISO 17025	months		
External - national arrangements (e.g. regulator)		months		
Internal		months		
Other		months		
None				


Box H: Declaration of Compliance

Do you have to make an annual declaration of compliance?

Who does this go to (e.g. regulator, certification body)?



Quality Audits





Audit Intervals

	Min.	Mode	Max.
External Formal (e.g. ISO 17025, ISO 9001 etc.)	12	12	36
External National (e.g. regulator)	1	12	72
Internal	1	12	36



Quality Audits

- There is wide acceptance of the importance of audit: 74% of services have internal audits, and two-thirds undergo formal QMS audits.
- The typical interval between external audits is 12 months.
- National regulatory requirements vary. Some audit very frequently, others infrequently.
- Internal practices also vary between IMSs.
- A substantial minority of IMSs are required to make an annual declaration of compliance.



QA in Dosimetry

Type Testing

- 91% of IMSs use dosemeters that have undergone formal or recognised type testing.
- 97% of IMSs use dosemeters that have undergone any type testing.
- the importance of type testing is widely accepted.

Traceability

- most services have established traceability.
- the typical interval between traceability exercises is 12 months.
- services should compare their practice with the RP160 recommendations.



Use of Proficiency Testing

